



## Polynomial interpolation for fractional iteration

### An interpolation-approach to fractional iteration of a powerseries-function

*Abstract: In this short article I discuss one way to approach a fractional iterate of a function  $f(x)$ , which is defined by a powerseries/exponential-series (this concept is not only applicable to the tetration (T-tetration, U-tetration)-problem). The fractional iterate is here understood as interpolation of the coefficients of the set of powerseries, which represent the consecutive integer iterates of the given function, to build a new powerseries using these interpolated coefficients. Two simple examples are given.*

*The article is simply an extended version of a discussion in the sci.math-newsgroup.*

---

*Update 3*

Assume a function, defined as powerseries in  $x$ :

$$(1) \quad f(x) = ax + bx^2 + cx^3 + dx^4 + \dots$$

Then the iterations  $f(f(x)), f(f(f(x))), \dots$  are expressible as new powerseries

$$(2) \quad f^{(1)}(x) = f(x) \quad f^{(2)}(x) = f(f(x)) \quad f^{(h+1)}(x) = f(f^{(h)}(x)) \quad f^{(0)}(x) = x$$

The new coefficients of, for instance of  $f^{(2)}(x)$  are finite combinations of the original coefficients  $a, b, c, \dots$  in the following way, which occur, if in (1) for  $x$  the function  $f(x)$  is inserted, the expressions expanded and like powers of  $x$  are collected:

$$(3) \quad \begin{aligned} f^{(2)}(x) &= a f(x) + b f(x)^2 + c f(x)^3 + \dots \\ &= a(ax + bx^2 + cx^3 + \dots) \\ &\quad + b(ax + bx^2 + cx^3 + \dots)(ax + bx^2 + cx^3 + \dots) \\ &\quad + c(ax + bx^2 + cx^3 + \dots)^3 \\ &\quad + \dots \\ &= a^2x + abx^2 + acx^3 + \dots \\ &\quad + a^2bx^2 + 2ab^2x^3 + \dots \\ &\quad + a^3cx^3 + \dots \\ &= a^2x + (ab+a^2b)x^2 + (ac + 2ab^2 + a^3c)x^3 + \dots \end{aligned}$$

This can be iterated. One may then observe the progression of the coefficients at each power of  $x$  along the iterations like

$$(4) \quad \begin{aligned} f^{(0)}(x) &= 1x \\ f^{(1)}(x) &= a x + b x^2 + c x^3 + \dots \\ f^{(2)}(x) &= a^2x + (ba^2+ba)x^2 + (ac + 2ab^2 + a^3c)x^3 + \dots \end{aligned}$$

when read column-wise.

If in our powerseries for  $f(x)$  the term  $a=1$  this simplifies significantly:

$$(5) \quad \begin{aligned} f^{(0)}(x) &= 1x \\ f^{(1)}(x) &= 1x + b x^2 + c x^3 + \dots \\ f^{(2)}(x) &= 1x + 2b x^2 + (2b^2 + 2c)x^3 + \dots \end{aligned}$$

and when this is written in a table we see, that – based on the progressions in the coefficients of each power of  $x$  – we can build polynomials in  $h$  to express the coefficients for each iterate of  $f^{(h)}(x)$  and to define continuous iterates as interpolations based on these coefficients-polynomials.

(6) Table of differences of coefficients along iterates of  $f^{(h)}(x)$ , using  $a=1$

diff. index	$f^{(0)}(x)$	$f^{(1)}(x)$	$f^{(2)}(x)$	$f^{(3)}(x)$
$\Delta^0$	$1x$ $0$ $0$ $0$ $0$ ...	$1x$ $1b x^2$ $1c x^3$ $1d x^4$ $1e x^5$ ...	$1x$ $(2b) x^2$ $(2c + 2bb) x^3$ $(2d + 5bc + b^3) x^4$ $(2e + 3(2bd + c^2) + 5b^2c) x^5$ ...	$1x$ $(3b) x^2$ $(3c + 6b^2) x^3$ $(3d + 15bc + 9b^3) x^4$ $(3e + 9(2bd + c^2) + 41b^2c + 10b^4) x^5$ ...
$\Delta^1$		$0x$ $1b x^2$ $1c x^3$ $1d x^4$ $1e x^5$	$0x$ $(1b) x^2$ $(1c + 2bb) x^3$ $(1d + 5bc + b^3) x^4$ $(1e + 3(2bd + c^2) + 5b^2c) x^5$	$0x$ $(1b) x^2$ $(1c + 4b^2) x^3$ $(1d + 10bc + 8b^3) x^4$ $(1e + 6(2bd + c^2) + 36b^2c + 10b^4) x^5$
$\Delta^2$			$0x$ $0x^2$ $(2bb) x^3$ $(5bc + b^3) x^4$ $(3(2bd + c^2) + 5b^2c) x^5$	$0x$ $0x^2$ $(2bb) x^3$ $(5bc + 7b^3) x^4$ $(3(2bd + c^2) + 31b^2c + 10b^4) x^5$
$\Delta^3$				$0x$ $0x^2$ $0x^3$ $(6b^3) x^4$ $(26b^2c + 10b^4) x^5$

Legend: linear progression quadratic progression cubic progression biquadratic progression of coefficients

**Example 1:**

If all coefficients  $a, b, c, d, \dots = 1$  we have the powerseries

$$(7) \quad f(x) = f^{(1)}(x) = 1x + 1x^2 + 1x^3 + \dots$$

the iterates

$$(8) \quad \begin{aligned} f^{(1)}(x) &= 1x + 1x^2 + 1x^3 + \dots \\ f^{(2)}(x) &= 1x + 2x^2 + 4x^3 + \dots \\ f^{(3)}(x) &= 1x + 3x^2 + 9x^3 + \dots \\ &\dots \\ f^{(h)}(x) &= 1x + hx^2 + h^2x^3 + \dots \end{aligned}$$

with the according ranges of convergence for  $x$  (not discussed here). The coefficients of the powerseries for the fractional iterates  $h$  are then simply the powers of  $h$ . (which, btw, is another access to describe the first column of the  $h$ 'th (continuous) powers of the pascalmatrix  $P$ ).

**Example 2:**

If the coefficients  $a, b, c, d, \dots$  follow the sequence of reciprocals of factorials, as it is with the second column in the matrix **S2** (factorial scaled Stirling-numbers 2'nd kind, see appendix) we get

$$(a=1, b=1/2!, c=1/3!, \dots)$$

$$(9) \quad \begin{aligned} f^{(1)}(x) &= 1x + 1/2 x^2 + 1/6 x^3 + \dots \\ f^{(2)}(x) &= 1x + 1 x^2 + 5/6 x^3 + 5/8 x^4 + 13/30 x^5 + \dots \\ f^{(3)}(x) &= 1x + 3/2 x^2 + 2 x^3 + 5/2 x^4 + 179/60 x^5 + \dots \\ &\dots \end{aligned}$$

which agrees with the matrix-computation of the powers of **S2**. The coefficients for the powerseries, which reflects the  $h$ 'th iterate (which may also be fractional) are then expressible as polynomials in  $h$ :

(10) *Polynomials for coefficients at  $k$ 'th power of  $x$*

$x^k$	<u>coefficients as functions of <math>h</math></u>
$x^{1*}$	(1 )
$x^{2*}$	(0 + 1/2 h )
$x^{3*}$	(0 - 1/12 h + 1/4 h <sup>2</sup> )
$x^{4*}$	(0 1/48 h - 5/48 h <sup>2</sup> + 1/8 h <sup>3</sup> )
...	...

So we get the powerseries for an arbitrary (fractional or complex)  $h$ 'th iterate:

$$(11) \quad f^{(h)}(x) = 1x + (1/2 h) x^2 + (1/4 h^2 - 1/12 h) x^3 + (1/8 h^3 - 5/48 h^2 + 1/48 h) x^4 + \dots$$

which defines an interpolated (continuous)  $h$ 'th iterate of the function  $f(x) = \exp(x) - 1$  (or: "U-tetration"  $U_e(x, h)$  of height  $h$ )

**Extension to infinite series**

We may assume U-tetration – series of consecutive heights,  $h=0,1,2,3,\dots$ , where  $U(x,1)=exp(x)-1$

$$AU(x) = x - U(x,1) + U(x,2) - U(x,3) + \dots$$

This is reflected by the infinite alternating series of  $h$  in formula (11), expressed by the (modified) Dirichlet-eta-function  $\eta(-k) = 0^k - 1^k + 2^k - \dots + \dots$ (start at index 0)

$$\begin{aligned}
 (11) \quad AU(x) &= 1x \\
 &- (1x + (1/2*1)x^2 + (1/4*1^2 - 1/12*1)x^3 + (1/8*1^3 - 5/48*1^2 + 1/48*1)x^4 + \dots) \\
 &+ (1x + (1/2*2)x^2 + (1/4*2^2 - 1/12*2)x^3 + (1/8*2^3 - 5/48*2^2 + 1/48*2)x^4 + \dots) \\
 &- (1x + (1/2*3)x^2 + (1/4*3^2 - 1/12*3)x^3 + (1/8*3^3 - 5/48*3^2 + 1/48*3)x^4 + \dots) \\
 &\dots \\
 &= \frac{\eta(0)x + (1/2*\eta(-1))x^2 + (1/4*\eta(-2) - 1/12*\eta(-1))x^3 + (1/8*\eta(-3) - 5/48*\eta(-2) + 1/48*\eta(-1))x^4 + \dots}{1/2x - 1/8x^2 + 1/48x^3 + 1/96x^4 - 19/1920x^5 - 13/7680x^6 + 2623/322560x^7 \dots}
 \end{aligned}$$

These coefficients occur similarly, if the geometric-series of  $S2$  is computed by the formula

$$AS2 = (I + S2)^{-1}$$

Unfortunately, the absolute values of coefficients grow after they approach a local minimum; the first 32 terms of the powerseries are

$$\begin{aligned}
 AU(x) = &0.5*x - 0.125*x^2 + 0.0208333*x^3 + 0.0104167*x^4 - 0.00989583*x^5 - 0.00169271*x^6 + 0.00813182*x^7 \\
 &- 0.00113157*x^8 - 0.0111201*x^9 + 0.00600352*x^{10} + 0.0232243*x^{11} - 0.0248572*x^{12} \\
 &- 0.0689663*x^{13} + 0.122171*x^{14} + 0.275591*x^{15} - 0.745013*x^{16} - 1.41855*x^{17} + 5.62062*x^{18} \\
 &+ 9.06553*x^{19} - 51.7967*x^{20} - 69.5330*x^{21} + 574.804*x^{22} + 617.490*x^{23} - 7577.81*x^{24} \\
 &- 6052.46*x^{25} + 117228.*x^{26} - 59786.4*x^{27} - 2.10483E6*x^{28} - 430171.*x^{29} + 4.34384E7*x^{30} \\
 &- 5.25132E6*x^{31} - 1.02150E9*x^{32} + O(x^{33})
 \end{aligned}$$

The growthrate of the absolute values of the coefficients of this powerseries is hypergeometric, so the radius of convergence is zero, and also cannot simply be summed via Euler-summation.

If we construct the series of negative heights instead of positive heights, we get the same coefficients, only of inverse sign, except at the first term, and this defines then

$$AL(x) = x - \log(1+x) + \log(1+\log(1+x)) - \dots + \dots$$

The sum of  $AU(x)$  and  $AL(x)$ , both expressed by the powerseries, is then

$$AU(x) + AL(x) = 0.5x + 0.5x = x$$

because all coefficients vanish; but as we have seen in a discussion in sci.math, this result disagrees with numerical computations, which were done by evaluation of the partial sums in some convergent cases (with appropriate bases  $b$ ). This discrepancy can still not be explained.

**Appendix:**

*Pari/GP (%-commands are Pari-TTY meta-tags redirecting matrix-output into separate GUI-windows)*

```
tmpser1=Ser([a,b,c,d,e,f])*x
tmpser2 =subst(tmpser1,x,tmpser1)
tmpser3 =subst(tmpser2,x,tmpser1)
tmpser4 =subst(tmpser3,x,tmpser1)
tmpser5 =subst(tmpser4,x,tmpser1)

%box >tst subst(Mat(polcoeffs(tmpser2))~,a,1)
```

*Powers of S2:*

$$\begin{bmatrix} 1 & . & . & . & . \\ 0 & 1 & . & . & . \\ 0 & 1/2 & 1 & . & . \\ 0 & 1/6 & 1 & 1 & . \\ 0 & 1/24 & 7/12 & 3/2 & 1 \\ 0 & 1/120 & 1/4 & 5/4 & 2 & 1 \end{bmatrix} S2 \begin{bmatrix} 1 & . & . & . & . \\ 0 & 1 & . & . & . \\ 0 & 1 & 1 & . & . \\ 0 & 5/6 & 2 & 1 & . \\ 0 & 5/8 & 8/3 & 3 & 1 \\ 0 & 13/30 & 35/12 & 11/2 & 4 & 1 \end{bmatrix} S2^2 \begin{bmatrix} 1 & . & . & . & . \\ 0 & 1 & . & . & . \\ 0 & 3/2 & 1 & . & . \\ 0 & 2 & 3 & 1 & . \\ 0 & 5/2 & 25/4 & 9/2 & 1 \\ 0 & 179/60 & 11 & 51/4 & 6 & 1 \end{bmatrix} S2^3$$

*tetration-Index*

<http://go.helms-net.de/math/tetdocs/index.htm>

*Gottfried Helms, 22.12.2007*